
Citation:

Helme, M and Low, C and Emmonds, S (2019) A Kinetic and Kinematic Analysis of The Rear Foot Elevated Split Squat 5RM Test. In: United Kingdom Strength and Conditioning Association National Conference 2019, 15 June 2019 - 16 June 2019, Stadium MK, Milton Keynes. (Unpublished)

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/6000/>

Document Version:

Conference or Workshop Item (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

A Kinetic and Kinematic Analysis of The Rear Foot Elevated Split Squat 5RM Test

Mark Helme, Dr Stacey Emmonds, Dr Chris Low¹
¹ Institute of Sport, Physical activity and Leisure, Carnegie School of Sport.
Mark.helme@leedsbeckett.ac.uk

Introduction

The rear foot elevated split squat (RFESS) is a multi-joint, unilateral resistance exercise, commonly used in strength and conditioning (McCurdy, 2017). McCurdy, Langford et al. (2004) and McCurdy and Langford (2005) have previously reported the RFESS as a reliable measure of unilateral leg strength (1RM ICC, 0.97- 0.99). The aim of this study was to firstly quantify the kinetic and kinematic characteristics of the RFESS 5RM test protocol. Secondly to profile the intra-set differences between repetitions.

Methods

26 volunteers were recruited, (age = 23.8 ±4.6 years, mass = 88.1 ±10.7kg, height = 1.79±0.1m), all subjects were engaged in a structured strength and conditioning program. Kinetic data was collected from the front and rear foot through two independent Kistler force plates. Kinematic data was captured through Qualysis Track Manager System at 250Hz (Qualysis AB, Gothenburg, Sweden) using 10 cameras (six ceiling mounted and four, floor mounted).







Data was extracted and exported to Biomechanics Toolbar, and a Butterworth, fourth order filter applied. Subsequent data was further exported to R for analysis. A second data set was created in Biomechanics Toolbar, by time normalising the data to 101 time points.

RESULTS

The mean load lifted was 84kg ±16.8kg (0.96 ±0.18 kg/kg). The mean vertical displacement of the bar was 0.38 ± 0.06m, mean concentric velocity was 0.32 ±0.05m/s and peak concentric velocity was 0.49 ±0.11m/s. The mean vertical ground reaction force (vGRF) of the lead foot was 1432.54±200.87N, (1.66 ±0.20BW). The lead foot produced 83.53±4.03% of total vGRF. There were unclear differences in all kinetic variables between all repetitions, except for peak (vGRF) of the lead foot only (1.90±0.28BW) of Repetition 5, which was very likely larger. Repetitions 1 and 2 were likely to very likely to have higher mean concentric velocities (MCV) than repetitions 4 and 5.

Stronger participants were able to achieve lower concentric velocities, both as a set mean and the 5th repetition of the set

Time normalisation to 101 time points of all repetitions found peak displacement occurred at point 51±6, where peak lead and rear foot vGRF occurred at points 56±33 and 60±40 respectively.



Conclusion:

The RFESS 5RM is valid and reliable method of measuring unilateral leg strength. Mean force contributions across the repetition of ≈15% with unreliable occurrences of peak rear foot vGRF, indicate a reliance on the lead for vertical displacement both eccentrically and concentrically. The RFESS can therefore be considered a unilateral exercise as no clear role of the rear foot could be defined.

A multi-repetition protocol can be used to determine maximal strength, yet intra-set differences may not exist prior to completion of the final repetition. It is recommended that the mean of the repetitions is taken when summarising the kinematic and kinetic variables in a multi repetition test. However, the final repetition MCV maybe used to determine relative intensity of the set or achievement of maximal performance. Based on the data from this study, this value may be 0.27m/s.